

Patterns of Motorbike Accidents Related Head Injuries in Patients Presenting to a Tertiary Care Hospital of Peshawar

MOHAMMAD ISHAQ, ADNAN AHMED, IKRAM ALAM

Musawer Khan, Syed Nasir Shah, Mumtaz Ali

Department of Neurosurgery, Lady Reading Hospital, Peshawar – Pakistan

ABSTRACT

Objective: To determine the different patterns of motorbike related head injuries in patients presenting to a tertiary care hospital of Peshawar.

Materials and Methods: It was a prospective (observational) study, which was conducted at the Neurotrauma ward of Neurosurgery department, Lady Reading Hospital, Peshawar from January 2016 to June 2016. Total number of patients were 246. We included those patients who were having impaired Glasgow Coma Scale (GCS), danger signs and having some abnormal findings on CT brain. We excluded those patients who were having GCS 15/15 and those patients of head injury having associated co-morbid conditions.

Results: There were 230 (93.49%) males and 16 (6.5%) females. Age range was 1 to 80 years. Majority of patients were drivers (55.28%). Only 6 (2.43%) drivers used helmet. Ninety patients (36.58%) had an ICU stay of more than one week. Seventy six (30.89%) patients were operated and 170 (69.1%) patients were treated conservatively. Mortality was 14.63% in our study.

Conclusion: Majority were males in our study, most of the patients who sustained head injury in motorbike accidents were drivers. Thirty six percent of the patients needed ICU stay of more than 1 week. Maximum of the patients needed conservative treatment.

Keywords: Road Traffic Accidents, Head Injury, Motorbike Accident.

Abbreviations: RTAs: Road Traffic Accidents. LOC: Loss of Consciousness. GOS: Glasgow Outcome Score.

INTRODUCTION

Road traffic accidents (RTAs) are the major cause of diseases and mortality throughout the world, particularly in the developing countries.¹⁻³ It is ranked ninth globally among major causes of disability and will be the third leading cause by 2020.⁴ It is estimated that more than 1 million people were killed in road traffic accidents and more than this were injured (approximately 300 deaths/ day).⁵ With increasing motorization in developing countries, it will be a major health problem in the near future.⁶⁻⁸ Injuries from RTAs results in major financial losses and productivity losses and leaves an astonishing effect on patients and their families. The effects of road traffic injuries are more in developing countries as compared to developed coun-

tries.^{9,10}

A study done in Singapore and Vietnam concluded higher proportion of motorbike related injuries of 49.1% and 62% respectively, while in Nigeria Madubuze and Labinjo reported 54% crash injuries.¹¹⁻¹⁴ In Pakistan road traffic accident related injuries are the 5th leading cause of loss of healthy life and second important cause of disability. In Pakistan, mortality rate from RTAs is 4 – 5/million population and 15/ million vehicles. Non lethal injuries are reported to be 200/100,000 population in major cities of Pakistan including Peshawar. There is a drastic increase in number of motor bikes across the province and especially in Peshawar. During the last decade the number of registered and unregistered vehicle increased up to

5 folds which is the major cause of congestion on the roads and ultimately leading to increase number of road traffic accidents. There is 14 fold increases in RTAs in Pakistan.¹⁵

Motor bikes need quick decisions as he has to respond quickly to stop or to turn a side in case of bad roads.^{16,17} Increase in the number of two wheel rides are associated with increased number of injuries and deaths. They are more prone to head and spine injuries. Most of the bike riders are young people and usually they tend to adopt risky attitudes and behaviors, they also do not use helmet due to which there is an increased risk of head injuries.¹⁸ The increasing risk of accidents and associated injuries depends on patients' age, passenger type, attitude and behavior of rider, use of helmet and road condition.¹⁹⁻²¹ The main reason behind increasing number of RTAs is due to increase in number of motor bikes, it's easy availability on installments, poor law enforcement, and non use of safety measures by the passengers and poor road infrastructure.^{21,22}

There is lack of published data on motorbike related injuries specially the head and spine injuries. The two departments involved in the recording of data are the police department and hospital. There is always discrepancy in publication of data by either of the department.²³⁻²⁵ Findings of our study will help in highlighting the issue and will enable the policy makers to formulate laws that will minimize mortalities and morbidities associated with motor bike injuries.

MATERIALS AND METHODS

A prospective study was done for duration of 6 months from January 2016 to June 2016 at neurotrauma ward of Neurosurgery department, Lady Reading Hospital, Peshawar. A total number of 246 patients were studied.

Inclusion Criteria

All those patients who were having impaired GCS with danger signs and having some abnormal findings on CT brain. Both genders were included irrespective of their age. Pedestrians were also included in the study.

Exclusion Criteria

All those patients who were having associated comorbid conditions, like diabetes, hypertension, ischemic heart disease and patients on anti-platelet drugs.

Protocol

All patients were examined by ATLS (Advanced trauma life support) protocol and were stabilized in the trauma room. After resuscitation a detailed history was obtained and a pre designed proforma was filled that included patients age, gender, passenger type, use of helmet, arrival GCS and other associated non neurological injuries.

After history and examination patient were subjected to investigations that included C.T scan brain with bone window, Digital X-ray cervical/ thoracic/lumbar spine followed by MRI or 3-D CT of the affected part of the spine.

Head injury was classified as mild, moderate and severe on the basis of GCS, GCS 13 – 15, 9 – 12, and less than or equal to 8 respectively. Danger signs like, post head injury amnesia, Loss of Consciousness (LOC) > 20 min, seizures and vomiting were also included. Patients with normal GCS having no danger signs and normal CT brain were discharged from neurotrauma without admission.

RESULTS

A total of 246 patients were included in the study. Results were analyzed using SPSS version 20. Males were 230 (93%) and females were 16 (7%). Age was divided in 8 groups (Table 1). Among the affectees the drivers were 136, back seaters were 72 and pedestrian were 38. Regarding the use of helmet 6 patients were having helmet, 200 were having no helmet and about 40 patients there is no idea of helmet use. Patients were referred from other hospitals as well and some came directly to the neurotrauma department (Table 2). Frequency of head injury type is shown in table 3.




Fig. 1: Gender Distribution.

Table 1: Age Distribution.

Age Group (Years)	Number of Patients	Percentage
1 – 10	22	8.94%
11 – 20	86	34.95%
21 – 30	54	21.95%
31 – 40	24	9.75%
41 – 50	18	7.31%
51 – 60	14	5.69%
61 – 70	4	1.62%
71 – 80	2	0.81%
Total	246	100%

Table 2: Patient's Address.

Location	Approximate Distance (KM) & time (min) From LRH, Peshawar	Frequency
Bajawar	173/192	6
Bannu	218/221	6
Bunir	134/165	12
Charsadda	31/32	18
Chitral	326/561	4
Dir (upper)	213/307	12
Hangu	112/150	4
Karak	141/156	16
Kohat	74/98	12
Khyber Agency	32/55	4
Malakand (Chakdara)	103/149	10
Mardan	60/60	46
Nowshehra	9/50	8
Swabi	98/72	6
Swat (Saidu Shareef)	141/212	12
Peshawar	5-10/30	74

CT scan findings are shown in Table 4. Patient with associated spine injuries were 16 and those having long bone injuries were 18. Thirty-six patients died in our study, 90 patients had an ICU stay of more than one week. Seventy-six patients were operated and 170 patients were treated conservatively. Glasgow outcome score is shown in Table 5.

Table 3: Frequency of Head Injury Type.

Type of Head Injury	GCS	Frequency
Mild	13 – 15	20.32% (50)
Moderate	9 – 12	37.39% (92)
Severe	3 – 8	42.27% (104)

Table 4: CT Scan Findings.

CT Brain (with Bone Window) Findings	Frequency
Depressed skull fractures	7.3% (18)
Subdural hematomas	13.82% (34)
Epidural hematomas	23.57% (58)
Contusions	25.60% (63)
Traumatic Subarachnoid hemorrhage	18.29% (45)
Pneumocephalus	13% (32)
Brain edema	48.70% (120)

Table 5: Glasgow Outcome Score.

Glasgow Outcome Score (GOS)	Frequency
1	12.19% (30)
2	5.28% (13)
3	12.60% (31)
4	24.39% (60)
5	45.52% (112)

DISCUSSION

Head injuries are the most common injuries related to motor bike injuries. Road traffic accident is considered as “silent epidemic” of the industrialized world.²⁶ Traumatic brain injuries are the hallmark of road traffic accidents specially related to motor bikers those

who are non-helmeted. Loss of consciousness and amnesia after trauma are the two main markers of severity of brain injury, but for ease it has been classified into three grades on the basis of GCS at presentation. Mild is the one having GCS 13 – 15, moderate having GCS 9 – 12 and severe having GCS 3 – 8. Patients with GCS more than 13 are debatable to be included in the traumatic brain injury sub group but studies have shown that there are increased chances of abnormal radiological findings in such people.^{27,28} In our study we also had a significant number (20.32%) of patients in the mild sub group who were having abnormal CT brain findings.

We found that males are the predominant victims of neurosurgical traumas and those in second and third decade are mostly affected.^{29,30} Regarding the age of the patient, the prevalence of traumatic brain injuries are more in two decades, so it represents a bimodal pattern. One peak is in the second and third decade and another in the 6th decade after 65 years of age, but in our study we found that the prevalence is more in second decade which is consistent with the previous study findings and it is followed by third decade and 4th decade. It represents that mostly the younger group of population which are from 11 years to 40 years are mostly affected.³¹ A study done in Karachi shows similar results with our studies. They also reported a high incidence in third decade of life. Likewise Jooma et al³² & Raja et al³³ conducted two different studies and concluded that second and fourth decade are mostly affected groups.

The average time duration to reach the hospital trauma unit was approximately 3 hours; most of the people were brought from far flung areas without any special ambulance (having no ventilator facility) and even in private vehicles, this result in the delay of delivering first aid services and urgent surgical intervention if needed.³⁴

Road traffic accidents are the most common cause of traumatic brain injuries which may be attributed towards traffic congestion, lack of traffic rules and regulations. There is an upsurge of motorization in Pakistan which has led to increased number of deaths due to road traffic accidents.³⁵ Naddumba³⁶ and Okeniyi et al,³⁷ reported that pedestrians are majority of victims affected and we found 15.44% pedestrians to be affected. It is due to the fact that majority of cities in the developing countries like Pakistan, pedestrian's signs are absent and if present are not followed by pedestrians and drivers. In Pakistan there is lack of knowledge and awareness regarding traffic rules and

regulations and most of bike riders do not use helmets during driving. We found that 81.30% didn't use helmets and regarding 16.30% there is no idea about helmet use. Our results are quite consistent with the study done in Tehran by Zarger et al³⁸ who found 91.4% of the cases were non helmeted. Another study showed 72% of the cases as non helmeted ones. Literature search shows severity of injury is high among non helmeted patients.^{39,40} A study shown only 3% of helmet use which is also evident in our study that only 2.43% of the cases had used helmets which is a critical issue and less than satisfactory, helmet use can result in prevention of TBI among bike riders.⁴¹⁻⁴⁴ Alcohol intoxication can also result in increasing prevalence of road traffic accidents. However, there is low prevalence of alcohol intoxication in our part of the world so only 0.8% of the cases were having history of alcohol ingestion. Similar study found 1.88% alcohol intoxicated patients.^{45,36}

CT scan is the choice of initial imaging modality during the first 24 hours following head injury.⁴⁶ we also performed CT scan brain in all cases in our study. CT is more accurate in detecting the hematomas and bony pathologies as compared to MRI.⁴⁷ In case of deteriorating of neurological status of the patient, CT scan should be done as found by Papa M et al.⁴⁸ and Stippler et al.⁴⁹ Another study proposed that CT should be done in cases of some predictors like headache, vomiting, loss of consciousness or amnesia and alcohol intoxication.^{50,51} Leong LB et al. found in a study done on 2038 patients, that CT scan should be done in the presence of LOC, vomiting and amnesia.⁵² We also did CT brain in the presence of any of these signs in addition to arrival GCS of the patient. We found abnormal findings in all of our cases. Those who were not having any positive CT scan findings were excluded from our study.

In the current study 7.3% of cases were having depressed skull fractures, 18.82% subdural hematomas, 23.57% epidural hematomas, 48.70% contusions, 18.29% traumatic sub arachnoid hemorrhages and 13% pneumocephalus. A study is done which shows findings like epidural hematomas in significant number of cases, 5% having subdural hematomas with high mortality rates. However, mortality is related to the arrival GCS and time interval between trauma and surgery.⁵³ A study shown 12.5% mortality related to traumatic brain injuries due to motorbike injuries.⁵⁴ Our study showed 12.19% mortality. A study of 344 patients in Nepal showed 4.5% mortality. Whereas Hitimana et al found 13.2% mortality in his study.⁵⁵

Intracranial hemorrhage, brain edema contusions and depressed skull fractures are the common CT findings following head trauma during road traffic crashes. A study showed 21.6% patients with brain edema.⁵⁶ We found 48.70% patients with brain edema (plus other pathologies) which is the second most common finding in our study after epidural hematomas. A study done which shows brain edema as the most common CT finding in patients with acute head injury.⁵⁷ We found 18.21% cases of traumatic sub arachnoid hemorrhage which is in contrast with the study which shows 41% of cases having traumatic subarachnoid hemorrhage.⁵⁸ The most common CT scan finding in a study conducted in Karachi, was brain contusion (14.1%); others included traumatic sub-arachnoid hemorrhage (7.1%), subdural hematoma (7.6%), extra-dural hematoma (5.8%) and depressed skull fracture (4.6%).³⁴

CONCLUSION

Majority were males in our study, most of the patients who sustained head injury in motorbike accidents were drivers. Thirty six percent of the patients needed ICU stay of more than 1 week. Maximum of the patients needed conservative treatment.

Address for correspondence:

Dr. Mohammad Ishaq

Registrar Neurosurgery A Department

Lady Reading Hospital, Peshawar – Pakistan

Cell No: 0300 5973012

Email: drmohammadishaq@yahoo.com

REFERENCES

1. Ameratunga S HM, Norton R. Road traffic injuries: confronting disparities to address a global health problem. *Lancet*, 2006; 367: 1533–40.
2. Mohan D. Road safety in less-motorized environments: future concerns. *Int. J. Epidemiol.* 2002; 31: 527–32.
3. Nantulya VM RM. The neglected epidemic: road traffic injuries in developing countries. *BMJ*. 2002; 324: 1139–41.
4. Peden M mK, Krug E. Injury: a leading cause of the global burden of disease, 2000. Geneva: WHO; 2002. The injury chart book 2000 (2002).
5. Peden MM, McGee K, Krug E, Injuries WHO, Dept VP. Injury: a leading cause of the global burden of disease, 2000: Dept. of Injuries and Violence Prevention, Noncommunicable Diseases and Mental Health Cluster, WHO; 2002.
6. Odero W GP, Zwi A. Road traffic injuries in developing countries: a comprehensive review of epidemiological studies. *Trop Med Int Health*, 1997; 2 (5): 445–60.
7. Jacobs G AA, Astrop A. Estimating global road fatalities. London: Transport Research Laboratory. 2000 (TRL Report 445).
8. Murray CJ LA. The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries and risk factors in 1990 and projected to 2020. (Global burden of disease and injury series). Boston: Harvard School of Public Health; 1996: 1.
9. Bazarian JJ, Blyth B, Mookerjee S, He H, McDermott MP: Sex differences in outcome after mild traumatic brain injury. *J Neurotrauma*, 2010; 27: 527–539.
10. Bell RS, Mossop CM, Dirks MS, Stephens FL, Mulligan L, Ecker R, Neal CJ, Kumar A, Tigno T, Armonda RA: Early decompressive craniectomy for severe penetrating and closed head injury during wartime. *Neurosurg Focus*, 2010; 28: E1.
11. Nantulya VM RM. Equity dimensions of road traffic injuries in low and middle-income countries. *Inj Control Saf Promot.* 2003; 10 (1-2): 13-20.
12. Madubueze CC, Chukwu CO, Omoke NI, Oyakhilome OP, Ozo C. Road traffic injuries as seen in a Nigerian teaching hospital. *Int Orthop.* Jul 11. Pub Med PMID: 20623283.
13. Tham KY, Seow E, Lau G. Pattern of injuries in helmeted motorcyclists in Singapore. *Emergency medicine journal: EMJ.* 2004 Jul; 21 (4): 478-82.
14. Labinjo M, Juillard C, Kobusingye OC, Hyder AA. The burden of road traffic injuries in Nigeria: results of a population-based survey. *Inj Prev.* 2009 Jun; 15 (3): 157-62.
15. Ghaffar A, Hyder A, Mastoor M, Shaikh I. Injuries in Pakistan: directions for future health policy. *Health policy and planning*, 1999; 14: 11.
16. Mayou R, Tyndel S, Bryant B. Long-term outcome of motor vehicle accident injury. *Am Psychosomatic Soc* 1997: 578-84.
17. Peden M, Scurfield R, Sleet D, Mohan D, Hyder AA, Jarawan E, et al. World report on road traffic injury prevention. World Health Organization Geneva; 2004.
18. Hyder AA, Morrow RH. Applying burden of disease methods to developing countries: case study from Pakistan. *Am J Public Health*, 2000; 90: 1235-40.
19. A, Rehman. The Burden of Road Traffic Injuries in South Asia: A Commentary. *JCPSP.* 2004; 14: 707-8.
20. Razzak JA, Luby SP. Estimating deaths and injuries due to road traffic accident in Karachi, Pakistan, through the capture-recapture method. *Int J Epidemiol.* 1998; 27: 866-70.
21. Hyder AA, Ghaffar A, M asood TI. Motor Vehicle crashes in Pakistan: the emerging epidemic. *Injury prevention*, 2006; 6: 199-202.
22. Sexton B, Baughan C, Elliott M, Maycock G. The accident risk of motorcyclists. Crowthorne: Transport Research Laboratory, 2004.

23. Kenneth S. O benski P. M otorcycle A ccident Recon-
struction and Litigation, 3rd Ed. Lawyers & Judges
Publishing, 2002.
24. Tham KY, Seow E, Lau G. Pattern of injuries in hel-
meted motorcyclists in Singapore. *Emerg Med J.* 2004;
478-82.
25. Skalkidou A, Petridou E, Papadopoulos FC, Dessypris
N, Trichopoulos D. Factors affecting motorcycle helmet
use in the population of Greater Athens, Greece. *Injury
Prevention*, 1999; 264-7.
26. larke DD, Ward P, Bartle C, Truman W. In-depth Study
of Motorcycle Accidents. Road Safety Research Report,
2004: 54.
27. Christian WJ, Carroll M, Meyer K, Vitaz TW, Franklin
GA. Motorcycle helmets and head injuries in Kentucky,
1995-2000. *J Ky Med Assoc.* 2003; 101: 21.
28. Van HT, S in g h asiv an o n P, K aew kungw al J, Suri-
yawongpaisal P, Khai LH. Estimation of non-fatal road
traffic injuries in Thai Nguyen, Vietnam using capture-
recapture method. *Southeast Asian J Trop Med Public
Health*, 2006; 37: 405.
29. Shukla D, Devi BI: Mild traumatic brain injuries in
adults. *J Neurosci Rural Pract.* 2010; 1 (2): 82-88.
30. Lee B, Newberg A: Neuroimaging in Traumatic Brain
Imaging. *Neuro Rx.* 2005; 2 (2): 372-83.
31. Servadei F, Teasdale G, Merry G: Neurotraumatology
Committee of the World Federation of Neurosurgical
Societies. Defining acute mild head injury in adults: A
proposal based on prognostic factors, diagnosis, and
management. *J Neurotrauma*, 2001; 18: 657–664.
32. Masson F et al.: Epidemiology of severe brain injuries:
A prospective population-based study. *J Trauma*, 2001;
51 (3): 481-489.
33. Stevens JA, Dellinger AM: Motor vehicle and fall re-
lated deaths among older Americans 1990–98: Sex, race,
and ethnic disparities. *Inj Prev.* 2002; 8 (4): 272-275.
34. Umerani MS¹, Abbas A, Sharif S. Traumatic brain inju-
ries: experience from a tertiary care centre in Pakistan.
Turk Neurosurg. 2014; 24 (1): 19-24.
35. Thurman D: Head Trauma: Basic, preclinical, and clini-
cal directions. In: Miller L, Ayes R (ed). *Head Trauma:
Basic, Preclinical, and Clinical Directions.* New York:
John Wiley and Sons, 2001: 327–347.
36. Jooma R, Ahmed S, Zarden AM: Comparison of Two
Surveys of head injured patients presenting during a
calendar year to an Urban Medical Centre 32 years
apart. *J Pak Med Ass.* 2005; 55 (12): 630-632.
37. Raja IA, Vohra AH, Ahmed M: Neurotrauma in Pakis-
tan. *World J Surg.* 2001; 25: 1230–1237.
38. Jinnah Postgraduate Medical Centre, Half Yearly Rep-
ort Jan to Jun & Half Yearly Comparison of 2008 to
2011. Road Traffic Injury Research & Prevention Cen-
tre; 2011 Aug.
39. Hyder AA, Ghaffar A, Masood TI: Motor vehicle cra-
shes in Pakistan: The emerging epidemic. *Inj Prev.*
2000; 6: 199-202.
40. Naddumba, E.K. A cross sectional retrospective study
of boda injuries at Mulago Hospital in Kampala, Uga-
nda. *ECAJS.* 2004; 9: 44-47.
41. Okeniyi, J.A. et al. Motorcycle injury: an emerging
menace to child health in Nigeria. *Internet J Pediatr
Neonatal.* 2005; 5 (1).
42. J. Mayrose, “The effects of a mandatory motorcycle
helmet law on helmet use and injury patterns among
motorcyclist fatalities,” *J Safety Res.* 2008; Vol. 39,
No. 4: pp. 429–432.
43. P. F. Umebese and S. U. Okukpo, “Motorcycle acci-
dents in a Nigeria University campus: a one year study
of pattern of trauma sustained in the University of Ben-
in,” *Niger J Clin Pract.* 2001; Vol. 10: pp. 33–36.
44. P. Siviroj, K. Peltzer, S. Pengpid, and S. Morarit, “Hel-
met use and associated factors among thai motorcyclists
during Songkran festival,” *Int J Environ Res Public
Health*, 2012; Vol. 9, No. 9: pp. 3286–3297.
45. B. A. Solagberu, C. K. P. Ofoegbu, A. A. Nasir, O. K.
Ogundipe, A.O. Adekanye, and L.O. Abdur-Rahman,
“Motorcycle injuries in a developing country and the
vulnerability of riders, passengers, and pedestrian,” *Inj
Prev.* 2006; Vol. 12, No. 4: pp. 266–268.
46. Raja IA, Vohra AH, Ahmed M: Neurotrauma in Pakis-
tan. *World J Surg.* 2001; 25: 1230–1237.
47. M. Richter, D. Otte, U. Lehmann et al., “Head injury
mechanisms in helmet-protected motorcyclists: pros-
pective multicenter study,” *J Trauma—Injury, Infection
and Critical Care*, 2001; Vol. 51, No. 5: pp. 949–958.
48. Kidwell CS et al: Comparison of MRI and CT for dete-
ction of acute intracerebral hemorrhage. *JAMA.* 2004;
292 (15): 1823-1830.
49. Papa L, et al: Performance of the Canadian CT Head
Rule and the New Orleans Criteria for predicting any
traumatic intracranial injury on computed tomography
in a United States Level I trauma center. *Acad Emerg
Med.* 2012; 19 (1): 2-10.
50. Stippler M, et al. Utility of routine follow-up head CT
scanning after mild traumatic brain injury: A systematic
review of the literature. *Emerg Med J.* 2012; 29 (7):
528-532.
51. Bainbridge J, Khirwadkar H, Hourihan MD: Vomiting -
is this a good indication for CT head scans in patients
with minor head injury? *Br J Radiol.* 2012; 85 (1010):
183-186.
52. Sharif-Alhoseini M, Khodadadi H, Chardoli M, Rahi-
mi-Movaghar V: Indications for brain computed tomo-
graphy scan after minor head injury. *J Emerg Trauma
Shock*, 2011; 4 (4): 472-476.
53. Leong LB, Sukarom S, Vasu A, Hian LG: Identifying
predictors of an abnormal computed tomographic scan
among patients with a head injury and a Glasgow Coma
Scale of 15. *Eur J Emerg Med Epub ahead of print*,
2012.
54. J. Hitimana, M. Perez, A. Kinasha, and I. Kakande,
“Clinical presentation and outcome of neurosurgical

- conditions at Butare teaching hospital, Rwanda,” East Central Afr J Surg. 2009; Vol. 14, No. 1: pp. 50–56.
55. Nolan S: Traumatic brain injury: A review. Crit Care Nurs Q. 2005; 28: 188-194.
56. Nnadi MON, Bankole OB, Fente BG. Motorcycle-Related Traumatic Brain Injuries: Helmet Use and Treatment Outcome. J Neurosci. 2015; 2015: 696-787. Doi:10.1155/2015/696787
57. Zimmerman, Robert A., et al. "Computed Tomography of Pediatric Head Trauma: Acute General Cerebral Swelling 1." Radiology, 1978; 126.2: 403-408.
58. Servadei, Franco, et al. "Traumatic subarachnoid hemorrhage: demographic and clinical study of 750 patients from the European brain injury consortium survey of head injuries." Neurosurgery, 2002. 50.2: 261-269.

AUTHORS DATA

Name	Post	Institution	E-mail	Role of Authors	
Dr. Mohammad Ishaq	Registrar	Department of Neurosurgery A, Lady Reading Hospital, Peshawar–Pakistan.	drmohammadishaq@yahoo.com	Paper Writing	
Dr. Adnan Ahmed					Data Collection
Dr. Ikram Alam					Results and Analysis
Dr. Musawer Khan					Research and Discussion
Dr. Syed Nasir Shah					Proof Reading
Dr. Mumtaz Ali	Professor				Overall Supervision

Date of Submission: 07-06-2017

Date of Printing: 22-06-2017

Peer Reviewed by Dr. Babar Butt and Chief Editor Prof. Dr. Muhammad Anwar Chaudary and others.